

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
21 July 2005 (21.07.2005)

PCT

(10) International Publication Number  
**WO 2005/066182 A1**

(51) International Patent Classification<sup>7</sup>: **C07D 495/04**

(74) Agents: **ASAMURA, Kiyoshi** et al.; Room 331, New Ohtemachi Bldg., 2-1, Ohtemachi 2-chome, Chiyoda-ku, Tokyo, 1000004 (JP).

(21) International Application Number:  
PCT/JP2005/000318

(22) International Filing Date: 6 January 2005 (06.01.2005)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
2004-001310 6 January 2004 (06.01.2004) JP

(71) Applicant (for all designated States except US): **TAISHO PHARMACEUTICAL CO., LTD.** [JP/JP]; 24-1, Takada 3-chome, Toshima-ku, Tokyo, 1708633 (JP).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **NAKAZATO, Atsuro** [JP/JP]; c/o Taisho Pharmaceutical Co., Ltd., 24-1, Takada 3-chome, Toshima-ku, Tokyo, 1708633 (JP). **OKUBO, Taketoshi** [JP/JP]; c/o Taisho Pharmaceutical Co., Ltd., 24-1, Takada 3-chome, Toshima-ku, Tokyo, 1708633 (JP). **NOZAWA, Dai** [JP/JP]; c/o Taisho Pharmaceutical Co., Ltd., 24-1, Takada 3-chome, Toshima-ku, Tokyo, 1708633 (JP). **TAMITA, Tomoko** [JP/JP]; c/o Taisho Pharmaceutical Co., Ltd., 24-1, Takada 3-chome, Toshima-ku, Tokyo, 1708633 (JP). **KENNIS, Ludo, E.J.** [BE/BE]; c/o Janssen Pharmaceutica N.V., Turnhoutseweg 30, Beerse, B-2340 (BE).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.


(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

**Published:**

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: THIENOPYRIMIDINE AND THIENOPYRIDINE DERIVATIVES SUBSTITUTED WITH CYCLIC AMINO GROUP



(57) Abstract: An object of the present invention is to provide an antagonist against CRF receptors which is effective as a therapeutic or prophylactic agent for diseases in which CRF is considered to be involved, such as depression, anxiety, Alzheimer's disease, Parkinson's disease, Huntington's chorea, eating disorder, hypertension, gastral diseases, drug dependence, epilepsy, cerebral infarction, cerebral ischemia, cerebral edema, cephalic external wound,

inflammation, immunity-related diseases, alpecia, irritable bowel syndrome, sleep disorders, dermatitides, schizophrenia, pain, etc. A thienopyrimidine or thienopyridine derivative substituted with a cyclic amino group represented by the following formula [I] has a high affinity for CRF receptors and is effective against diseases in which CRF is considered to be involved.

## DESCRIPTION

THIENOPYRIMIDINE AND THIENOPYRIDINE DERIVATIVES  
SUBSTITUTED WITH CYCLIC AMINO GROUP

## DETAILED DESCRIPTION OF THE INVENTION

## TECHNICAL FIELD

The present invention relates to a therapeutic agent for diseases in which corticotropin releasing factor (CRF) is considered to be involved, such as depression, anxiety, Alzheimer's disease, Parkinson's disease, Huntington's chorea, eating disorder, hypertension, gastral diseases, drug dependence, cerebral infarction, cerebral ischemia, cerebral edema, cephalic external wound, inflammation, immunity-related diseases, alpecia, irritable bowel syndrome, sleep disorders, epilepsy, dermatitides, schizophrenia, pain, etc.

10

## DESCRIPTION OF THE PRIOR ART

CRF is a hormone comprising 41 amino acids (Science, 213, 1394-1397, 1981; and J. Neurosci., 7, 88-100, 1987), and it is suggested that CRF plays a core role in biological reactions against stresses (Cell. Mol. Neurobiol., 14, 579-588, 1994; Endocrinol., 132, 723-728, 1994; and Neuroendocrinol. 61, 445-452, 1995). For CRF, there are the following two paths: a path by which CRF acts on peripheral immune system or sympathetic nervous system through hypothalamus-pituitary-adrenal system, and a path by which CRF functions as a neurotransmitter in central nervous system (in Corticotropin Releasing Factor: Basic and Clinical Studies of a Neuropeptide, pp. 29-52, 1990). Intraventricular administration of CRF to hypophysectomized rats and normal rats causes an anxiety-like symptom in both types of rats (Pharmacol. Rev., 43, 425-473, 1991; and Brain Res. Rev., 15, 71-100, 1990). That is, there are suggested the participation of CRF in hypothalamus-pituitary-adrenal system and the pathway by which CRF functions as a neurotransmitter in central nervous system.

25

The review by Owens and Nemeroff in 1991 summarizes diseases in which CRF is involved (Pharmacol. Rev., 43, 425-474, 1991). That is, CRF is

involved in depression, anxiety, Alzheimer's disease, Parkinson's disease, Huntington's chorea, eating disorder, hypertension, gastrointestinal diseases, drug dependence, inflammation, immunity-related diseases, etc. It has recently been reported that CRF is involved also in epilepsy, cerebral infarction, cerebral  
5 ischemia, cerebral edema, and cephalic external wound (Brain Res. 545, 339-342, 1991; Ann. Neurol. 31, 48-498, 1992; Dev. Brain Res. 91, 245-251, 1996; and Brain Res. 744, 166-170, 1997). Accordingly, antagonists against CRF receptors are useful as therapeutic agents for the diseases described above.

WO02/002549, WO97/29110 and WO98/47903 disclose thienopyridine  
10 and thienopyrimidine derivatives respectively as CRF receptor antagonists. However, none disclose the compounds provided in the present invention.

#### PROBLEM(S) TO BE SOLVED BY INVENTION

An object of the present invention is to provide an antagonist against CRF  
15 receptors which is effective as a therapeutic or prophylactic agent for diseases in which CRF is considered to be involved, such as depression, anxiety, Alzheimer's disease, Parkinson's disease, Huntington's chorea, eating disorder, hypertension, gastral diseases, drug dependence, epilepsy, cerebral infarction, cerebral ischemia, cerebral edema, cephalic external wound, inflammation, immunity-related diseases,  
20 alpecia, irritable bowel syndrome, sleep disorders, dermatitides, schizophrenia, pain, etc.


#### MEANS FOR SOLVING PROBLEM

The present inventors earnestly investigated thienopyrimidine or  
25 thienopyridine derivatives substituted with a cyclic amino group that have a high affinity for CRF receptors, whereby the present invention has been accomplished.

The present invention is thienopyrimidine or thienopyridine derivatives substituted with a cyclic amino group explained below.


A thienopyrimidine or thienopyridine derivative substituted with a cyclic  
30 amino group represented by the following formula [I]:

3



5

(wherein the cyclic amino group is represented by the following formula [II]):



10

in which the cyclic amino group is a 3- to 8-membered saturated cyclic amine or a 3- to 8-membered saturated cyclic amine bridged with C<sub>1-5</sub>alkylene or C<sub>1-4</sub>alkylene-O-C<sub>1-4</sub>alkylene between any different two carbon atoms of the cyclic amine, which cyclic amine is substituted with a group represented by -(CR<sup>1</sup>R<sup>2</sup>)<sub>m</sub>-(CHR<sup>3</sup>)<sub>n</sub>-X, R<sup>4</sup> and R<sup>5</sup> independently on the same or different carbon atoms of the cyclic amine;

15

X is cyano, hydroxy, -CO<sub>2</sub>R<sup>8</sup> or -CONR<sup>9</sup>R<sup>10</sup>;

Y is N or CR<sup>11</sup>;

20

R<sup>1</sup> is hydrogen, hydroxy, C<sub>1-5</sub>alkyl, C<sub>1-5</sub>alkoxy-C<sub>1-5</sub>alkyl or hydroxy-C<sub>1-5</sub>alkyl;

R<sup>2</sup> is hydrogen or C<sub>1-5</sub>alkyl;

R<sup>3</sup> is hydrogen, cyano, C<sub>1-5</sub>alkyl, C<sub>1-5</sub>alkoxy-C<sub>1-5</sub>alkyl or hydroxy-C<sub>1-5</sub>alkyl;

m is an integer selected from 0, 1, 2, 3, 4 and 5;

25

n is 0 or 1;

R<sup>4</sup> is hydrogen, hydroxy, hydroxy-C<sub>1-5</sub>alkyl, cyano, cyano-C<sub>1-5</sub>alkyl or C<sub>1-5</sub>alkyl;

R<sup>5</sup> is hydrogen or C<sub>1-5</sub>alkyl;

30

R<sup>6</sup> is hydrogen, C<sub>1-5</sub>alkyl, C<sub>3-8</sub>cycloalkyl, C<sub>3-8</sub>cycloalkyl-C<sub>1-5</sub>alkyl, hydroxy, C<sub>1-5</sub>alkoxy, C<sub>3-8</sub>cycloalkyloxy, halogen, C<sub>1-5</sub>alkylthio or -N(R<sup>12</sup>)R<sup>13</sup>;

R<sup>7</sup> is hydrogen, halogen, C<sub>1-5</sub>alkyl, C<sub>3-8</sub>cycloalkyl, C<sub>3-8</sub>cycloalkyl-C<sub>1-5</sub>alkyl, hydroxy, C<sub>1-5</sub>alkoxy, C<sub>3-8</sub>cycloalkyloxy, -N(R<sup>14</sup>)R<sup>15</sup>, -CO<sub>2</sub>R<sup>16</sup>, -CON(R<sup>17</sup>)R<sup>18</sup>, cyano, nitro, C<sub>1-5</sub>alkylthio, trifluoromethyl or trifluoromethoxy;

Ar is aryl or heteroaryl which aryl or heteroaryl is unsubstituted or substituted with 1 or more substituents, which are the same or different, selected from the group consisting of halogen, C<sub>1-5</sub>alkyl, C<sub>3-8</sub>cycloalkyl, C<sub>2-5</sub>alkenyl, C<sub>2-5</sub>alkynyl, C<sub>1-5</sub>alkoxy, C<sub>1-5</sub>alkylthio, C<sub>1-5</sub>alkylsulfinyl, C<sub>1-5</sub>alkylsulfonyl, cyano, nitro, hydroxy, -CO<sub>2</sub>R<sup>19</sup>, -C(=O)R<sup>20</sup>, -CONR<sup>21</sup>R<sup>22</sup>, -OC(=O)R<sup>23</sup>, -NR<sup>24</sup>CO<sub>2</sub>R<sup>25</sup>, -S(=O)<sub>r</sub>NR<sup>26</sup>R<sup>27</sup>, trifluoromethyl, trifluoromethoxy, difluoromethoxy, fluoromethoxy, methylenedioxy, ethylenedioxy and -N(R<sup>28</sup>)R<sup>29</sup>;

R<sup>8</sup> is hydrogen, C<sub>1-10</sub>alkyl, C<sub>3-8</sub>cycloalkyl, C<sub>3-8</sub>cycloalkyl-C<sub>1-5</sub>alkyl, aryl or aryl-C<sub>1-5</sub>alkyl;

R<sup>9</sup> and R<sup>10</sup> are the same or different, and independently are hydrogen, C<sub>1-5</sub>alkyl, C<sub>3-8</sub>cycloalkyl, C<sub>3-8</sub>cycloalkyl-C<sub>1-5</sub>alkyl, aryl or aryl-C<sub>1-5</sub>alkyl; or R<sup>9</sup> and R<sup>10</sup> form a ring selected from saturated 3 to 8 membered ring with the attached nitrogen atom, wherein one of the carbon atoms of such saturated 3 to 8 membered ring is optionally replaced by an oxygen or sulfur atom or by N-Z wherein Z is hydrogen, benzyl or C<sub>1-5</sub>alkyl;

R<sup>11</sup> is hydrogen, halogen or C<sub>1-5</sub>alkyl;

R<sup>12</sup>, R<sup>13</sup>, R<sup>14</sup> and R<sup>15</sup> are the same or different, and independently are hydrogen or C<sub>1-5</sub>alkyl;

R<sup>16</sup>, R<sup>19</sup> and R<sup>25</sup> are the same or different, and independently are hydrogen or C<sub>1-5</sub>alkyl, C<sub>3-8</sub>cycloalkyl, C<sub>3-8</sub>cycloalkyl-C<sub>1-5</sub>alkyl, aryl or aryl-C<sub>1-5</sub>alkyl;

R<sup>17</sup>, R<sup>18</sup>, R<sup>20</sup>, R<sup>21</sup>, R<sup>22</sup>, R<sup>23</sup>, R<sup>24</sup>, R<sup>26</sup>, R<sup>27</sup>, R<sup>28</sup> and R<sup>29</sup> are the same or different, and independently are hydrogen, C<sub>1-5</sub>alkyl or C<sub>3-8</sub>cycloalkyl;

r is 1 or 2)

, individual isomers thereof or racemic or non-racemic mixtures of isomers thereof, pharmaceutically acceptable prodrugs thereof or pharmaceutically acceptable salts and hydrates thereof.

The terms used in the present specification have the following meanings.

The term "a 3- to 8-membered saturated cyclic amine" means aziridine, azetidine, pyrrolidine, piperidine, azepane or azocane.

The term "C<sub>1-5</sub>alkylene" means a straight or branched chain alkylene of 1 to 5 carbon atoms, such as methylene, ethylene, propylene, trimethylene, tetramethylene, pentamethylene or the like.

The term "a 3- to 8-membered saturated cyclic amine bridged with C<sub>1-</sub>

salkylene or C<sub>1-4</sub>alkylene-O-C<sub>1-4</sub>alkylene between any different two carbon atoms of the cyclic amine" includes, for example, 8-azabicyclo[3.2.1]oct-8-yl, 9-azabicyclo[3.3.1]non-9-yl, 7-azabicyclo[2.2.1]hept-7-yl, 3-oxa-7-azabicyclo[3.3.1]non-7-yl and 3-oxa-9-azabicyclo[3.3.1]non-9-yl.

5           The term "C<sub>1-5</sub>alkyl" means a straight chain or branched chain alkyl group of 1 to 5 carbon atoms, such as methyl, ethyl, propyl, isopropyl, butyl, isobutyl, *t*-butyl, *sec*-butyl, pentyl, isopentyl or the like.

          The term "C<sub>1-5</sub>alkoxy" means a straight chain or branched chain alkoxy group of 1 to 5 carbon atoms, such as methoxy, ethoxy, propoxy, isopropoxy, butoxy, isobutyloxy, pentyloxy, isopentyloxy or the like.

          The term "C<sub>1-5</sub>alkoxy-C<sub>1-5</sub>alkyl" means a substituted C<sub>1-5</sub>alkyl group having the above-mentioned C<sub>1-5</sub>alkoxy group as the substituent, such as methoxymethyl, 2-methoxyethyl, 2-ethoxyethyl or the like.

          The term "hydroxy-C<sub>1-5</sub>alkyl" means a substituted C<sub>1-5</sub>alkyl group having hydroxy group, such as hydroxymethyl, 1-hydroxyethyl, 2-hydroxyethyl, 1-hydroxypropyl, 2-hydroxypropyl, 3-hydroxypropyl, 4-hydroxybutyl, 5-hydroxypentyl or the like.

          The term "cyano-C<sub>1-5</sub>alkyl" means a substituted C<sub>1-5</sub>alkyl group having cyano group, such as cyanomethyl, 1-cyanoethyl, 2-cyanoethyl, 3-cyanopropyl, 4-cyanobutyl, 5-cyanopentyl or the like.

          The term "C<sub>3-8</sub>cycloalkyl" means a cyclic alkyl group of 3 to 8 carbon atoms, such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl or the like.

          The term "C<sub>3-8</sub>cycloalkyl-C<sub>1-5</sub>alkyl" means a substituted C<sub>1-5</sub>alkyl group having the above-mentioned C<sub>3-8</sub>cycloalkyl as the substituent, such as cyclopropylmethyl, 2-cyclopropylethyl, 2-cyclopentylethyl or the like.

          The term "C<sub>3-8</sub>cycloalkyloxy" means a cyclic alkoxy group of 3 to 8 carbon atoms, such as cyclopropyloxy, cyclobutyloxy, cyclopentyloxy or the like.

          The term "halogen" means fluorine, chlorine, bromine or iodine atom.

30           The term "C<sub>1-5</sub>alkylthio" means a straight chain or branched chain alkylthio group of 1 to 5 carbon atoms, such as methylthio, ethylthio, propylthio or the like.

          The term "C<sub>1-5</sub>alkylsulfinyl" means a straight chain or branched chain

alkylsulfinyl group of 1 to 5 carbon atoms, such as methylsulfinyl, ethylsulfinyl, propylsulfinyl or the like.

The term "C<sub>1-5</sub>alkylsulfonyl" means a straight chain or branched chain alkylsulfonyl group of 1 to 5 carbon atoms, such as methylsulfonyl, ethylsulfonyl, propylsulfonyl or the like.

The term "aryl" means a monocyclic or bicyclic group of 6 to 12 ring carbon atoms having at least one aromatic ring, such as phenyl, naphthyl, or the like.

The term "heteroaryl" means a monocyclic or bicyclic group of 5 to 12 ring atoms having at least one aromatic ring having in its ring 1 to 4 atoms which may be the same or different and are selected from nitrogen, oxygen and sulfur, such as pyridyl, pyrimidinyl, imidazolyl, quinolyl, indolyl, benzofuranyl, quinoxaliny, benzo[1,2,5]thiadiazolyl, benzo[1,2,5]oxadiazolyl or the like.

The term "ary-C<sub>1-5</sub>alkyl" means a substituted C<sub>1-5</sub>alkyl group having the above-mentioned aryl as the substituent, such as benzyl, phenethyl or the like.

The term "C<sub>2-5</sub>alkenyl" means a straight chain or branched chain alkenyl group of 2 to 5 carbon atoms, such as vinyl, isopropenyl, allyl or the like.

The term "C<sub>2-5</sub>alkynyl" means a straight chain or branched chain alkynyl group of 2 to 5 carbon atoms, such as ethynyl, prop-1-ynyl, prop-2-ynyl or the like.

The phrase "aryl or heteroaryl which aryl or heteroaryl is unsubstituted or substituted with 1 or more substituents, which are the same or different, selected from the group consisting of halogen, C<sub>1-5</sub>alkyl, C<sub>3-8</sub>cycloalkyl, C<sub>2-5</sub>alkenyl, C<sub>2-5</sub>alkynyl, C<sub>1-5</sub>alkoxy, C<sub>1-5</sub>alkylthio, C<sub>1-5</sub>alkylsulfinyl, C<sub>1-5</sub>alkylsulfonyl, cyano, nitro, hydroxy, -CO<sub>2</sub>R<sup>19</sup>, -C(=O)R<sup>20</sup>, -CONR<sup>21</sup>R<sup>22</sup>, -OC(=O)R<sup>23</sup>, -NR<sup>24</sup>CO<sub>2</sub>R<sup>25</sup>, -S(=O)<sub>r</sub>NR<sup>26</sup>R<sup>27</sup>, trifluoromethyl, trifluoromethoxy, difluoromethoxy, fluoromethoxy, methylenedioxy, ethylenedioxy and -N(R<sup>28</sup>)R<sup>29</sup>" includes, for example, 2,4-dimethylphenyl, 2,6-dimethylphenyl, 2,4-dibromophenyl, 2-bromo-4-isopropylphenyl, 2,4-dichlorophenyl, 2,6-dichlorophenyl, 2-chloro-4-trifluoromethylphenyl, 4-methoxy-2-methylphenyl, 2-chloro-4-trifluoromethoxyphenyl, 4-isopropyl-2-methylthiophenyl, 2,4,6-trimethylphenyl, 4-bromo-2,6-dimethylphenyl, 4-bromo-2,6-diethylphenyl, 4-chloro-2,6-dimethylphenyl, 2,4,6-tribromophenyl, 2,4,5-tribromophenyl, 2,4,6-trichlorophenyl, 2,4,5-trichlorophenyl, 4-bromo-2,6-dichlorophenyl, 6-chloro-2,4-dibromophenyl,

2,4-dibromo-6-fluorophenyl, 2,4-dibromo-6-methylphenyl, 2,4-dibromo-6-methoxyphenyl, 2,4-dibromo-6-methylthiophenyl, 2,6-dibromo-4-isopropylphenyl, 2,6-dibromo-4-trifluoromethylphenyl, 2-bromo-4-trifluoromethylphenyl, 4-bromo-2-chlorophenyl, 2-bromo-4-chlorophenyl, 4-bromo-2-methylphenyl, 4-chloro-2-methylphenyl, 2,4-dimethoxyphenyl, 2,6-dimethyl-4-methoxyphenyl, 4-chloro-2,6-dibromophenyl, 4-bromo-2,6-difluorophenyl, 2,6-dichloro-4-trifluoromethylphenyl, 2,6-dichloro-4-trifluoromethoxyphenyl, 2,6-dibromo-4-trifluoromethoxyphenyl, 2-chloro-4,6-dimethylphenyl, 2-bromo-4,6-dimethoxyphenyl, 2-bromo-4-isopropyl-6-methoxyphenyl, 2,4-dimethoxy-6-methylphenyl, 6-dimethylamino-4-methylpyridin-3-yl, 2-chloro-6-trifluoromethylpyridin-3-yl, 2-chloro-6-trifluoromethoxypyridin-3-yl, 2-chloro-6-methoxypyridin-3-yl, 6-methoxy-2-trifluoromethylpyridin-3-yl, 2-chloro-6-difluoromethylpyridin-3-yl, 6-methoxy-2-methylpyridin-3-yl, 2,6-dimethoxypyridin-3-yl, 4,6-dimethyl-2-trifluoromethylpyrimidin-5-yl or 2-dimethylamino-6-methylpyridin-3-yl.

The “pharmaceutically acceptable salts” in the present invention include, for example, salts with an inorganic acid such as sulfuric acid, hydrochloric acid, hydrobromic acid, phosphoric acid, nitric acid or the like; salts with an organic acid such as acetic acid, oxalic acid, lactic acid, tartaric acid, fumaric acid, maleic acid, citric acid, benzenesulfonic acid, methanesulfonic acid, *p*-toluenesulfonic acid, benzoic acid, camphorsulfonic acid, ethanesulfonic acid, glucoheptonic acid, gluconic acid, glutamic acid, glycolic acid, malic acid, malonic acid, mandelic acid, galactaric acid, naphthalene-2-sulfonic acid or the like; salts with one or more metal ions such as lithium ion, sodium ion, potassium ion, calcium ion, magnesium ion, zinc ion, aluminium ion or the like; salts with amines such as ammonia, arginine, lysine, piperazine, choline, diethylamine, 4-phenylcyclohexylamine, 2-aminoethanol, benzathine or the like.


Prodrugs are also included in this invention. The term “prodrug” means a compound which is converted within the body, e.g. by hydrolysis in the blood, into its active form that has medical effects. The “pharmaceutically acceptable prodrugs” are described in, for example, Advanced Drug Delivery Reviews (1996) 19 (2) 115-130 and Tetrahedron Letter (2002) 43 1161-1164. The “pharmaceutically acceptable prodrugs thereof” in the present invention include, for example, esters such as methyl esters, ethyl esters, and the like when X is



carboxylic acid.

A compound of the present invention includes any isomers such as diastereomers, enantiomers, geometric isomers and tautomeric forms. In a compound represented by formula [I], if the cyclic amino group has one or more  
 5 chiral carbons and/or if there is an axial chirality between Ar and thienopyrimidine (or thienopyridine) ring, several stereoisomers (diastereomers or enantiomers) can exist. The compound of the present invention includes the individual isomers and the racemic and non-racemic mixtures of the isomers.


10 Preferable examples of the compound of the present invention are as follows.



That is preferable are compounds of the formula [III] in which X, m, n, the cyclic amino group, R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup> and Ar are as defined in above formula [I]. More preferable are compounds of the formula [III] in which X is  
 20 cyano; the cyclic amino group is a 4- to 7-membered saturated cyclic amine; n is 0; m is 0, 1, 2 and 3; R<sup>1</sup>, R<sup>2</sup>, R<sup>4</sup> and R<sup>5</sup> are hydrogen; R<sup>6</sup> is C<sub>1-5</sub>alkyl; R<sup>7</sup> is hydrogen or C<sub>1-5</sub>alkyl; and Ar is phenyl which phenyl is substituted with two or three substituents, which are the same or different, selected from the group consisting of halogen, C<sub>1-3</sub>alkyl, C<sub>1-3</sub>alkoxy, C<sub>1-3</sub>alkylthio, trifluoromethyl, trifluoromethoxy and  
 25 -N(R<sup>28</sup>)R<sup>29</sup> (wherein R<sup>28</sup> and R<sup>29</sup> are the same or different, and independently are hydrogen or C<sub>1-3</sub>alkyl). More preferable are compounds of the formula [III] in which X is cyano; the cyclic amino group is a 6-membered saturated cyclic amine; n is 0; m is 0 or 1; R<sup>1</sup>, R<sup>2</sup>, R<sup>4</sup> and R<sup>5</sup> are hydrogen; R<sup>6</sup> is C<sub>1-5</sub>alkyl; R<sup>7</sup> is hydrogen or C<sub>1-5</sub>alkyl; and Ar is phenyl which phenyl is substituted with two or three  
 30 substituents, which are the same or different, selected from the group consisting of halogen and C<sub>1-3</sub>alkyl.

Other preferable are compounds of the formula [III] in which X is hydroxy; the cyclic amino group is a 4- to 7-membered saturated cyclic amine; n is

0; m is an integer selected from 1, 2 and 3;  $R^1$ ,  $R^2$ ,  $R^4$  and  $R^5$  are hydrogen;  $R^6$  is  $C_{1-5}$ alkyl;  $R^7$  is hydrogen or  $C_{1-5}$ alkyl; and Ar is phenyl which phenyl is substituted with two or three substituents, which are the same or different, selected from the group consisting of halogen,  $C_{1-3}$ alkyl,  $C_{1-3}$ alkoxy,  $C_{1-3}$ alkylthio, trifluoromethyl, trifluoromethoxy and  $-N(R^{28})R^{29}$  (wherein  $R^{28}$  and  $R^{29}$  are the same or different, and independently are hydrogen or  $C_{1-3}$ alkyl). More preferable are compounds of the formula [III] in which X is hydroxy; the cyclic amino group is a 6-membered saturated cyclic amine; n is 0; m is an integer selected from 1, 2 and 3;  $R^1$ ,  $R^2$ ,  $R^4$  and  $R^5$  are hydrogen;  $R^6$  is  $C_{1-5}$ alkyl;  $R^7$  is hydrogen or  $C_{1-5}$ alkyl; and Ar is phenyl which phenyl is substituted with two or three substituents, which are the same or different, selected from the group consisting of halogen and  $C_{1-3}$ alkyl.



Other preferable are compounds of the formula [IV] in which X, m, n, the cyclic amino group,  $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^4$ ,  $R^5$ ,  $R^6$ ,  $R^7$ ,  $R^{11}$  and Ar are as defined in above formula [I]. More preferable are compounds of the formula [IV] in which X is cyano; the cyclic amino group is a 4- to 7-membered saturated cyclic amine; n is 0; m is 0 or 1;  $R^1$ ,  $R^2$ ,  $R^4$  and  $R^5$  are hydrogen;  $R^6$  is  $C_{1-5}$ alkyl;  $R^7$  is hydrogen or  $C_{1-5}$ alkyl;  $R^{11}$  is hydrogen or  $C_{1-5}$ alkyl; and Ar is phenyl which phenyl is substituted with two or three substituents, which are the same or different, selected from the group consisting of halogen,  $C_{1-3}$ alkyl,  $C_{1-3}$ alkoxy,  $C_{1-3}$ alkylthio, trifluoromethyl, trifluoromethoxy and  $-N(R^{28})R^{29}$  (wherein  $R^{28}$  and  $R^{29}$  are the same or different, and independently are hydrogen or  $C_{1-3}$ alkyl). More preferable are compounds of the formula [IV] in which X is cyano; the cyclic amino group is a 6-membered saturated cyclic amine; n is 0; m is 0 or 1;  $R^1$ ,  $R^2$ ,  $R^4$  and  $R^5$  are hydrogen;  $R^6$  is  $C_{1-5}$ alkyl;  $R^7$  is hydrogen or  $C_{1-5}$ alkyl;  $R^{11}$  is hydrogen; and Ar is phenyl which phenyl is substituted with two or three substituents, which are the same or different, selected from the group consisting of halogen and  $C_{1-3}$ alkyl.

Other preferable are compounds of the formula [IV] in which X is

hydroxy; the cyclic amino group is a 4- to 7-membered saturated cyclic amine; n is 0; m is an integer selected from 1, 2 and 3;  $R^1$ ,  $R^2$ ,  $R^4$  and  $R^5$  are hydrogen;  $R^6$  is  $C_{1-5}$ alkyl;  $R^7$  is hydrogen or  $C_{1-5}$ alkyl;  $R^{11}$  is hydrogen or  $C_{1-5}$ alkyl; and Ar is phenyl which phenyl is substituted with two or three substituents, which are the same or different, selected from the group consisting of halogen,  $C_{1-3}$ alkyl,  $C_{1-3}$ alkoxy,  $C_{1-3}$ alkylthio, trifluoromethyl, trifluoromethoxy and  $-N(R^{28})R^{29}$  (wherein  $R^{28}$  and  $R^{29}$  are the same or different, and independently are hydrogen or  $C_{1-3}$ alkyl). More preferable are compounds of the formula [IV] in which X is hydroxy; the cyclic amino group is a 6-membered saturated cyclic amine; n is 0; m is an integer selected from 1, 2 and 3;  $R^1$ ,  $R^2$ ,  $R^4$  and  $R^5$  are hydrogen;  $R^6$  is  $C_{1-5}$ alkyl;  $R^7$  is hydrogen or  $C_{1-5}$ alkyl;  $R^{11}$  is hydrogen; and Ar is phenyl which phenyl is substituted with two or three substituents, which are the same or different, selected from the group consisting of halogen and  $C_{1-3}$ alkyl.

The preferable cyclic amino group is a 6-membered saturated cyclic amine.

The preferable  $R^1$  is hydrogen.

The preferable  $R^2$  is hydrogen.

The preferable  $R^3$  is hydrogen.

The preferable  $R^4$  is hydrogen.

The preferable  $R^5$  is hydrogen.

The preferable  $R^6$  is  $C_{1-3}$ alkyl. The more preferable  $R^6$  is methyl.

The preferable  $R^7$  is hydrogen or  $C_{1-3}$ alkyl.

The preferable  $R^{11}$  is hydrogen.


The preferable Ar is phenyl which phenyl is substituted with two or three substituents, which are the same or different, selected from the group consisting of halogen,  $C_{1-3}$ alkyl,  $C_{1-3}$ alkoxy,  $C_{1-3}$ alkylthio, trifluoromethyl, trifluoromethoxy and  $-N(R^{28})R^{29}$  (wherein  $R^{28}$  and  $R^{29}$  are the same or different, and independently are hydrogen or  $C_{1-3}$ alkyl). The more preferable Ar is phenyl which phenyl is substituted with two or three substituents, which are the same or different, selected from the group consisting of halogen and  $C_{1-3}$ alkyl.

Especially preferable compounds of the present invention are:

{1-[7-(4-bromo-2,6-dimethyl-phenyl)-2-methyl-thieno[3,2-d]pyrimidin-4-yl]-piperidin-4-yl}-methanol


11

5




, {1-[7-(4-bromo-2,6-dimethyl-phenyl)-2,6-dimethyl-thieno[3,2-d]pyrimidin-4-yl]-piperidin-4-yl}-methanol

10




15 , 2-[1-[7-(4-bromo-2,6-dimethyl-phenyl)-2,6-dimethyl-thieno[3,2-d]pyrimidin-4-yl]-piperidin-4-yl]-ethanol

20



, {1-[7-(4-bromo-2,6-dimethyl-phenyl)-2,6-dimethyl-thieno[3,2-d]pyrimidin-4-yl]-piperidin-4-yl}-acetonitrile


25



30

, {1-[3-(2,4-dichloro-phenyl)-5-methyl-thieno[3,2-b]pyridin-7-yl]-piperidin-4-yl}-methanol


12



5


, {1-[5-methyl-3-(2,4,6-trimethyl-phenyl)-thieno[3,2-b]pyridin-7-yl]-piperidin-4-yl}-methanol

10



, {1-[3-(4-bromo-2,6-dimethyl-phenyl)-5-methyl-thieno[3,2-b]pyridin-7-yl]-piperidin-4-yl}-methanol


15



20

, {1-[3-(4-bromo-2,6-dimethyl-phenyl)-2,5-dimethyl-thieno[3,2-b]pyridin-7-yl]-piperidin-4-yl}-methanol


25



, {1-[3-(2,4-dibromo-phenyl)-5-methyl-thieno[3,2-b]pyridin-7-yl]-piperidin-4-yl}-methanol

30


13



5


, {1-[5-methyl-3-(2,4,6-trichloro-phenyl)-thieno[3,2-b]pyridin-7-yl]-piperidin-4-yl}-methanol

10



, 2-{1-[3-(4-bromo-2,6-dimethyl-phenyl)-5-methyl-thieno[3,2-b]pyridin-7-yl]-piperidin-4-yl}-ethanol


15



20


, 2-{1-[3-(4-bromo-2,6-dimethyl-phenyl)-2,5-dimethyl-thieno[3,2-b]pyridin-7-yl]-piperidin-4-yl}-ethanol

25



, 2-{1-[3-(2,4-dibromo-phenyl)-5-methyl-thieno[3,2-b]pyridin-7-yl]-piperidin-4-yl}-ethanol


30



14


, 2-{1-[5-methyl-3-(2,4,6-trichloro-phenyl)-thieno[3,2-b]pyridin-7-yl]-piperidin-4-yl}-ethanol

5




, 1-[5-methyl-3-(2,4,6-trimethyl-phenyl)-thieno[3,2-b]pyridin-7-yl]-piperidine-3-carbonitrile

10




15 , {1-[3-(4-bromo-2,6-dimethyl-phenyl)-5-methyl-thieno[3,2-b]pyridin-7-yl]-piperidin-4-yl}-acetonitrile

20




, {1-[3-(4-bromo-2,6-dimethyl-phenyl)-2,5-dimethyl-thieno[3,2-b]pyridin-7-yl]-piperidin-4-yl}-acetonitrile

25



30 , {1-[3-(2,4-dibromo-phenyl)-5-methyl-thieno[3,2-b]pyridin-7-yl]-piperidin-4-yl}-acetonitrile


15



5

and {1-[5-methyl-3-(2,4,6-trichloro-phenyl)-thieno[3,2-b]pyridin-7-yl]-piperidin-4-yl}-acetonitrile.

10




15

The compound of the formula [I] can be produced, for example, by the process shown in the following reaction scheme 1 (in the following reaction scheme,  $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^4$ ,  $R^5$ ,  $R^6$ ,  $R^7$ ,  $R^9$ ,  $R^{10}$ ,  $m$ ,  $n$ ,  $X$ ,  $Y$  and  $Ar$  are as defined above,  $LG$  is chloro, bromo, iodo, methanesulfonyloxy, benzenesulfonyloxy, toluenesulfonyloxy or trifluoromethanesulfonyloxy,  $X^a$  is carboxy, carbamoyl or  $-CO_2(C_{1-5}alkyl)$ ,  $X^b$  is  $CO_2(C_{1-5}alkyl)$  or  $CONR^9R^{10}$ ).

## 20 Reaction Scheme 1

25



## 30 Step 1:


Compound (3), a compound of the present invention, can be obtained by reacting Compound (1) with Compound (2) in an inert solvent in the presence or absence of a base. Herein, the base includes, for example, amines such as



triethylamine, *N,N*-diisopropylethylamine, pyridine and the like; inorganic bases such as sodium carbonate, potassium carbonate, sodium hydrogencarbonate, potassium hydrogencarbonate, sodium hydroxide, potassium hydroxide, barium hydroxide, sodium hydride and the like; metal alcoholates such as sodium methoxide, sodium ethoxide, potassium *tert*-butoxide and the like; metal amides such as sodium amide, lithium diisopropylamide and the like; and Grignard reagents such as methylmagnesium bromide and the like. The inert solvent includes, for example, alcohols such as methanol, ethanol, isopropyl alcohol, ethylene glycol and the like; ethers such as diethyl ether, tetrahydrofuran, 1,4-dioxane, 1,2-dimethoxyethane and the like; hydrocarbons such as benzene, toluene, xylene and the like; amides such as *N,N*-dimethylformamide, *N*-methylpyrrolidone, *N,N*-dimethylacetamide and the like; acetonitrile; dimethyl sulfoxide; pyridine; water; and mixtures of solvents selected from these inert solvents.

The compound of the present invention can be converted to a salt in an inert solvent with an inorganic acid such as sulfuric acid, hydrochloric acid, hydrobromic acid, phosphoric acid, nitric acid or the like, with an organic acid such as acetic acid, oxalic acid, lactic acid, tartaric acid, fumaric acid, maleic acid, citric acid, benzenesulfonic acid, methanesulfonic acid, *p*-toluenesulfonic acid, benzoic acid, camphorsulfonic acid, ethanesulfonic acid, glucoheptonic acid, gluconic acid, glutamic acid, glycolic acid, malic acid, malonic acid, mandelic acid, galactaric acid, naphthalene-2-sulfonic acid or the like, with an inorganic base such as lithium hydroxide, sodium hydroxide, potassium hydroxide, calcium hydroxide, magnesium hydroxide, zinc hydroxide, aluminium hydroxide or the like or with an organic base such as ammonia, arginine, lysine, piperazine, choline, diethylamine, 4-phenylcyclohexylamine, 2-aminoethanol, benzathine or the like. The inert solvent includes, for example, alcohols such as methanol, ethanol, isopropyl alcohol, ethylene glycol and the like; ethers such as diethyl ether, tetrahydrofuran, 1,4-dioxane, 1,2-dimethoxyethane and the like; esters such as ethyl acetate, ethyl formate and the like; ketones such as acetone, methylethylketone and the like; hydrocarbons such as benzene, toluene and the like; amides such as *N,N*-dimethylformamide, *N*-methylpyrrolidone, *N,N*-dimethylacetamide and the like; acetonitrile; dichloromethane; chloroform; dimethyl sulfoxide; pyridine; water; and mixtures of solvents selected from these inert solvents.


## Reaction Scheme 2



## Step 2:

The cyano group in compound (4), which was synthesized by the similar method described in step 1, can be converted to the carboxyl group, a C<sub>1</sub>-alkoxycarbonyl group or the carbamoyl group by using an acid or a base in an inert solvent or without any solvent. An oxidizing agent and/or a crown ether may be used as an additive in this reaction. Herein, the acid includes, for example, organic acids such as formic acid, acetic acid, trifluoroacetic acid, benzenesulfonic acid, methanesulfonic acid, *p*-toluenesulfonic acid, benzoic acid, trifluoromethanesulfonic acid and the like; inorganic acids such as sulfuric acid, hydrochloric acid, hydrobromic acid, phosphoric acid, polyphosphoric acid, nitric acid, boron trifluoride or the like. The base includes, for example, inorganic bases such as sodium carbonate, potassium carbonate, sodium hydrogencarbonate, potassium hydrogencarbonate, sodium hydroxide, potassium hydroxide, lithium hydroxide, barium hydroxide and the like; metal alcoholates such as sodium methoxide, sodium ethoxide, potassium *tert*-butoxide and the like. The oxidizing agent includes, for example, hydrogen peroxide, oxygen gas, manganese oxide and the like. The crown ether includes, for example, 18-crown-6, 15-crown-5 and the like. The inert solvent includes, for example, alcohols such as methanol, ethanol, isopropyl alcohol, ethylene glycol, *tert*-butanol and the like; ethers such as diethyl ether, tetrahydrofuran, 1,4-dioxane, 1,2-dimethoxyethane and the like; hydrocarbons such as benzene, toluene, xylene and the like; esters such as ethyl acetate, ethyl formate and the like; amides such as *N,N*-dimethylformamide, *N*-methylpyrrolidone, *N,N*-dimethylacetamide and the like; acetonitrile; dimethyl sulfoxide; pyridine; chloroform; dichloromethane; water; and mixtures of solvents selected from these inert solvents.

## Reaction Scheme 3



## Step 3:

Compound (7), a compound of the present invention, can be synthesized from Compound (6) by conventional methods for amidating a carboxyl group, esterification of a carboxyl group in the presence or absence of an acid or a base or alkylation of a carboxyl group with an alkylating reagent in an inert solvent. Herein, conventional methods for amidating a carboxyl group or esterification of a carboxyl group are: for example, the reaction *via* a mixed acid anhydride obtained by the reaction of Compound (6) with haloformic acid ester (e.g., ethyl chloroformate or isobutyl chloroformate) or an acid chloride (e.g., benzoyl chloride or pivaloyl chloride); the reaction in the presence of a condensing agent such as N,N'-dicyclohexylcarbodiimide (DCC), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (EDCI), carbonyldiimidazole (CDI), diphenylphosphoryl azide (DPPA), diethyl cyanophosphate or the like, optionally an additive such as 1-hydroxybenzotriazole (HOBt), N-hydroxysuccinimide, 4-dimethylaminopyridine or the like; or the reaction *via* an acid halide obtained by the reaction of Compound (6) with a halogenating reagent such as thionyl chloride, oxalyl chloride, or the like. The alkylating reagent is, for example, alkyl halide such as iodomethane, iodoethane, bromomethane, bromoethane and the like. The base includes amines such as triethylamine, N,N-diisopropylethylamine, pyridine, 1,8-diazabicyclo[5.4.0]undec-7-ene, 4-(dimethylamino)pyridine and the like; inorganic bases such as sodium carbonate, potassium carbonate, sodium hydrogencarbonate, potassium hydrogencarbonate and the like. The acid includes, for example, organic acids such as formic acid, acetic acid, trifluoroacetic acid, benzenesulfonic acid, methanesulfonic acid, *p*-toluenesulfonic acid, benzoic acid, trifluoromethanesulfonic acid and the like; inorganic acids such as sulfuric acid, hydrochloric acid, hydrobromic acid, phosphoric acid, polyphosphoric acid, nitric

acid or the like. The inert solvent includes, for example, alcohols such as methanol, ethanol, isopropyl alcohol, ethylene glycol and the like; ethers such as diethyl ether, tetrahydrofuran, 1,4-dioxane, 1,2-dimethoxyethane and the like; esters such as ethyl acetate, ethyl formate and the like; hydrocarbons such as benzene, toluene and the like; amides such as N,N-dimethylformamide, N-methylpyrrolidone, N,N-dimethylacetamide and the like; acetonitrile; dimethyl sulfoxide; pyridine; chloroform; dichloromethane; water; and mixtures of solvents selected from these inert solvents.

The compound of the present invention is useful as a therapeutic or prophylactic agent for diseases in which CRF is considered to be involved. For this purpose, the compound of the present invention can be formulated into tablets, pills, capsules, granules, powders, solutions, emulsions, suspensions, injections and the like by a conventional preparation technique by adding conventional fillers, binders, disintegrators, pH-adjusting agents, solvents, etc.


The compound of the present invention can be administered to an adult patient in a dose of 0.1 to 500 mg per day in one portion or several portions orally or parenterally. The dose can be properly increased or decreased depending on the kind of a disease and the age, body weight and symptom of a patient.

## 20 ENBODIMENTS OF THE INVENTION

The present invention is concretely explained with reference to the following examples and test example, but is not limited thereto.

### Example 1

Synthesis of {1-[7-(4-bromo-2,6-dimethyl-phenyl)-2,6-dimethyl-thieno[3,2-d]pyrimidin-4-yl]-piperidin-4-yl}-methanol hydrochloride (compound 1-004)



(1) A mixture of 7-(4-bromo-2,6-dimethyl-phenyl)-4-chloro-2,6-dimethyl-thieno[3,2-d]pyrimidine (500 mg), piperidin-4-ylmethanol (226 mg), *N,N*-


diisopropylethylamine (253 mg) in ethanol (1.5 mL) was heated at reflux for 1 day. The reaction mixture was cooled to room temperature, poured into a saturated aqueous sodium hydrogencarbonate, and then extracted with EtOAc. The organic layer was washed with brine, dried over anhydrous sodium sulfate and filtered. The filtrate was concentrated under reduced pressure and purified by a silica gel column chromatography (silica gel: Wako Gel (C200), eluent: hexane/EtOAc = 3 : 1) to obtain {1-[7-(4-bromo-2,6-dimethyl-phenyl)-2,6-dimethyl-thieno[3,2-d]pyrimidin-4-yl]-piperidin-4-yl}-methanol as a white solid (568 mg).

(2) To a suspension of {1-[7-(4-bromo-2,6-dimethyl-phenyl)-2,6-dimethyl-thieno[3,2-d]pyrimidin-4-yl]-piperidin-4-yl}-methanol (568 mg) in a mixture (1 : 1) of EtOH and EtOAc (2 mL) was added 4 M HCl in EtOAc (0.37 mL) under ice-cooling. The mixture was stirred overnight to afford a white crystal. The crystal was collected by filtration to give the title compound (532 mg).

Table 1 lists the compound obtained in Example 1 and compounds obtained by the similar procedure as described in Example 1.

### Example 2

{1-[7-(4-Bromo-2,6-dimethyl-phenyl)-2,6-dimethyl-thieno[3,2-d]pyrimidin-4-yl]-piperidin-4-yl}-acetic acid



(1) A mixture of {1-[7-(4-bromo-2,6-dimethyl-phenyl)-2,6-dimethyl-thieno[3,2-d]pyrimidin-4-yl]-piperidin-4-yl}-acetonitrile (350 mg) and KOH (492 mg) in a mixture of EtOH (1.5 mL) and H<sub>2</sub>O (1.0 mL) in a sealed tube was heated at 105 °C for 3 hours. After concentration of the reaction mixture under reduced pressure, 5 % KHSO<sub>4</sub> aqueous solution was added and extracted with CHCl<sub>3</sub>. The organic layer was washed with brine, dried over anhydrous Na<sub>2</sub>SO<sub>4</sub> and concentrated under reduced pressure. The residue was purified by a silica gel

21


column chromatography (silica gel: Wako Gel (C200), eluent:  $\text{CHCl}_3$ /methanol = 20 : 1) to obtain the title compound (164 mg).

Table 1 lists the compound obtained in Example 2 and compounds  
5 obtained by the similar procedure as described in Example 2.

### Example 3

2-{1-[7-(4-Bromo-2,6-dimethyl-phenyl)-2,6-dimethyl-thieno[3,2-  
d]pyrimidin-4-yl]-piperidin-4-yl}-acetamide

10




15

{1-[7-(4-Bromo-2,6-dimethyl-phenyl)-2,6-dimethyl-thieno[3,2-  
d]pyrimidin-4-yl]-piperidin-4-yl}-acetonitrile hydrochloride (30 mg) was dissolved  
in c  $\text{H}_2\text{SO}_4$  (0.5 mL) and the solution was stirred at room temperature for 20 hours.  
After addition of ice, the reaction mixture was made to alkaline (pH 7) with an  
aqueous NaOH solution and an aqueous  $\text{NaHCO}_3$  solution. The mixture was  
20 extracted with EtOAc and the organic layer was washed with brine, dried over  
anhydrous  $\text{Na}_2\text{SO}_4$  and concentrated under reduced pressure. The residue was  
purified by a silica gel column chromatography (silica gel: Wako Gel (C200),  
eluent: EtOAc) to obtain the title compound (20 mg) as a white crystal.

### 25 Example 4

{1-[7-(4-Bromo-2,6-dimethyl-phenyl)-2,6-dimethyl-thieno[3,2-  
d]pyrimidin-4-yl]-piperidin-4-yl}-acetic acid ethyl ester

30




A mixture of {1-[7-(4-bromo-2,6-dimethyl-phenyl)-2,6-dimethyl-

22

thieno[3,2-d]pyrimidin-4-yl]-piperidin-4-yl}-acetic acid (30 mg), iodoethane (97 mg) and  $K_2CO_3$  (17 mg) in DMF (1 mL) was stirred at room temperature for 16 hours. To the reaction mixture were added  $H_2O$  and EtOAc and separated. The organic layer was washed brine, dried over anhydrous  $Na_2SO_4$  and concentrated under reduced pressure. The residue was purified by a silica gel column chromatography (silica gel: Wako Gel (C200), eluent: Hexane/EtOAc = 4 : 1) to obtain the title compound (22 mg) as a white crystal.

























Table 1<sup>\*1</sup>























5



Com. No.	Ex. No.	$X-(CR^2R^3)_n-(CHR^1)_m$	Y	R <sup>6</sup>	R <sup>7</sup>	-Ar	melting point (°C) (solvent for crystallization)
1-001	1		N	CH <sub>3</sub>	H		amorphous
1-002	1		N	CH <sub>3</sub>	H		amorphous
1-003	1		N	CH <sub>3</sub>	H		177-180 <sup>*2</sup> (EtOAc/EtOH)
1-004	1		N	CH <sub>3</sub>	CH <sub>3</sub>		242-244 <sup>*2</sup> (EtOAc/EtOH)
1-005	1		N	CH <sub>3</sub>	H		192-194 <sup>*2</sup> (EtOH)
1-006	1		N	CH <sub>3</sub>	CH <sub>3</sub>		192-193 <sup>*2</sup> (EtOAc/EtOH)
1-007	1		N	CH <sub>3</sub>	H		amorphous
1-008	1		N	CH <sub>3</sub>	H		amorphous
1-009	1		N	CH <sub>3</sub>	H		163-165 <sup>*2</sup> (EtOAc/EtOH)



1-010	1		N	CH <sub>3</sub>	CH <sub>3</sub>		202-204* <sup>2</sup> (EtOAc/EtOH)
1-011	1		CH	CH <sub>3</sub>	H		amorphous
1-012	1		CH	CH <sub>3</sub>	H		amorphous
1-013	1		CH	CH <sub>3</sub>	H		amorphous
1-014	1		CH	CH <sub>3</sub>	H		228-230* <sup>2</sup> (EtOAc/EtOH)
1-015	1		CH	CH <sub>3</sub>	CH <sub>3</sub>		234-236* <sup>2</sup> (EtOAc/EtOH)
1-016	1		CH	CH <sub>3</sub>	H		196-199* <sup>2</sup> (EtOAc/EtOH)
1-017	1		CH	CH <sub>3</sub>	H		231-233* <sup>2</sup> (EtOAc/EtOH)
1-018	1		CH	CH <sub>3</sub>	H		172-174* <sup>2</sup> (EtOAc)
1-019	1		CH	CH <sub>3</sub>	CH <sub>3</sub>		182-184* <sup>2</sup> (EtOAc/EtOH)
1-020	1		CH	CH <sub>3</sub>	H		166-168* <sup>2</sup> (EtOAc/EtOH)
1-021	1		CH	CH <sub>3</sub>	H		158-160* <sup>2</sup> (EtOAc/EtOH)

1-022	1		CH	CH <sub>3</sub>	H		amorphous
1-023	1		CH	CH <sub>3</sub>	H		amorphous
1-024	1		CH	CH <sub>3</sub>	H		amorphous
1-025	1		CH	CH <sub>3</sub>	H		186-188 <sup>*2</sup> (EtOAc/IPE)
1-026	1		CH	CH <sub>3</sub>	CH <sub>3</sub>		135-137 <sup>*2</sup> (EtOAc/EtOH)
1-027	1		CH	CH <sub>3</sub>	H		179-182 <sup>*2</sup> (EtOAc/EtOH)
1-028	1		CH	CH <sub>3</sub>	H		203-205 <sup>*2</sup> (EtOAc)
1-029	2		N	CH <sub>3</sub>	CH <sub>3</sub>		268-270 (EtOAc)
1-030	2		N	CH <sub>3</sub>	CH <sub>3</sub>		222-224 (EtOAc)
1-031	3		N	CH <sub>3</sub>	CH <sub>3</sub>		212-124 <sup>*3</sup>
1-032	4		N	CH <sub>3</sub>	CH <sub>3</sub>		110-112 <sup>*3</sup>

\*1: Com. No. = compound number, Ex. No. = example number, solvent for crystallization; EtOAc = ethyl acetate, EtOH = ethanol, IPE = diisopropylether, Et = ethyl

Analytical data of non-crystal compounds are described below.

1-001:

MS (Pos, ES): 408 (M + 1)<sup>+</sup>, 410 (M + 3)<sup>+</sup>, 430 (M + Na)<sup>+</sup>, 432 (M + Na + 2)<sup>+</sup>;

NMR (300 MHz, CDCl<sub>3</sub>) δ 1.50-2.13 (5 H, m), 2.56 (3H, s), 3.48-3.62 (2H, m),

5 3.71-4.00 (3 H, m), 4.06-4.29 (2 H, m), 7.35 (1 H, dd, *J* = 2.0, 8.4 Hz), 7.52 (1 H, d, *J* = 2.0 Hz), 7.57 (1H, d, *J* = 8.4 Hz), 7.84 (1 H, s)

1-002:

MS (Pos, ES): 408 (M + 1)<sup>+</sup>, 410 (M + 3)<sup>+</sup>; HPLC Retention time: 9.69 (Xterra

10 MS C18 (Waters, Milford, MA) 3.5 μm, 4.6 x 100 mm); Flow rate 1.6 ml/min.

Three mobile phases (mobile phase A 95% 25mM ammonium acetate + 5% acetonitrile; mobile phase B: acetonitrile; mobile phase C: methanol) were employed to run a gradient condition from 100 % A to 50% B and 50% C in 10 min., to 100 % B in 1 min, 100% B for 3 min. and reequilibrate with 100 % A for

15 2.5 min)

1-007:

MS (Pos, ES): 436 (M + 1)<sup>+</sup>, 438 (M + 3)<sup>+</sup>; HPLC Retention time: 9.97 (Xterra

MS C18 (Waters, Milford, MA) 3.5 μm, 4.6 x 100 mm); Flow rate 1.6 ml/min.

20 Three mobile phases (mobile phase A 95% 25mM ammonium acetate + 5% acetonitrile; mobile phase B: acetonitrile; mobile phase C: methanol) were employed to run a gradient condition from 100 % A to 50% B and 50% C in 10 min., to 100 % B in 1 min, 100% B for 3 min. and reequilibrate with 100 % A for 2.5 min)

25

1-008:

MS (Pos, ES): 403 (M + 1)<sup>+</sup>, 405 (M + 3)<sup>+</sup>; HPLC Retention time: 9.94 (Xterra MS C18 (Waters, Milford, MA) 3.5 μm, 4.6 x 100 mm); Flow rate 1.6 ml/min.

30 Three mobile phases (mobile phase A 95% 25mM ammonium acetate + 5% acetonitrile; mobile phase B: acetonitrile; mobile phase C: methanol) were employed to run a gradient condition from 100 % A to 50% B and 50% C in 10 min., to 100 % B in 1 min, 100% B for 3 min. and reequilibrate with 100 % A for 2.5 min)

1-011:

MS (Pos, ES): 407 (M + 1)<sup>+</sup>, 409 (M + 3)<sup>+</sup>, 429 (M + Na)<sup>+</sup>, 431 (M + Na + 2)<sup>+</sup>;

NMR (300 MHz, CDCl<sub>3</sub>) δ 1.20-2.12 (5 H, m), 2.57 (3H, s), 2.80-3.06 (2H, m),

5 3.52-4.00 (5 H, m), 6.61 (1 H, s), 7.33 (1 H, dd, *J* = 2.0, 8.4 Hz), 7.51 (1 H, d, *J* = 2.0 Hz), 7.63 (1H, d, *J* = 8.4 Hz), 7.73 (1 H, s)

1-012:

MS (Pos, ES): 407 (M + 1)<sup>+</sup>, 409 (M + 3)<sup>+</sup>; **HPLC** Retention time: 10.02 (Xterra

10 MS C18 (Waters, Milford, MA) 3.5 μm, 4.6 x 100 mm); Flow rate 1.6 ml/min.

Three mobile phases (mobile phase A 95% 25mM ammonium acetate + 5% acetonitrile; mobile phase B: acetonitrile; mobile phase C: methanol) were employed to run a gradient condition from 100 % A to 50% B and 50% C in 10 min., to 100 % B in 1 min, 100% B for 3 min. and reequilibrate with 100 % A for

15 2.5 min)

1-013:

MS (Pos, ES): 381 (M + 1)<sup>+</sup>; **HPLC** Retention time: 9.22 (Xterra MS C18 (Waters, Milford, MA) 3.5 μm, 4.6 x 100 mm); Flow rate 1.6 ml/min. Three mobile phases

20 (mobile phase A 95% 25mM ammonium acetate + 5% acetonitrile; mobile phase B: acetonitrile; mobile phase C: methanol) were employed to run a gradient condition from 100 % A to 50% B and 50% C in 10 min., to 100 % B in 1 min, 100% B for 3 min. and reequilibrate with 100 % A for 2.5 min)

25 1-022:

MS (Pos, ES): 409 (M + 1)<sup>+</sup>; **HPLC** Retention time: 9.89 (Xterra MS C18 (Waters, Milford, MA) 3.5 μm, 4.6 x 100 mm); Flow rate 1.6 ml/min. Three mobile phases

(mobile phase A 95% 25mM ammonium acetate + 5% acetonitrile; mobile phase B: acetonitrile; mobile phase C: methanol) were employed to run a gradient  
30 condition from 100 % A to 50% B and 50% C in 10 min., to 100 % B in 1 min, 100% B for 3 min. and reequilibrate with 100 % A for 2.5 min)

1-023:

28

**MS** (Pos, ES): 402 ( $M + 1$ )<sup>+</sup>, 404 ( $M + 3$ )<sup>+</sup>; **HPLC** Retention time: 6.40 (Xterra MS C18 (Waters, Milford, MA) 3.5  $\mu$ m, 4.6 x 100 mm); Flow rate 1.6 ml/min. Three mobile phases (mobile phase A 95% 25mM ammonium acetate + 5% acetonitrile; mobile phase B: acetonitrile; mobile phase C: methanol) were employed to run a gradient condition from 100 % A to 50% B and 50% C in 10 min., to 100 % B in 1 min, 100% B for 3 min. and reequilibrate with 100 % A for 2.5 min)

1-024:

**MS** (Pos, ES): 376 ( $M + 1$ )<sup>+</sup>; **HPLC** Retention time: 6.21 (Xterra MS C18 (Waters, Milford, MA) 3.5  $\mu$ m, 4.6 x 100 mm); Flow rate 1.6 ml/min. Three mobile phases (mobile phase A 95% 25mM ammonium acetate + 5% acetonitrile; mobile phase B: acetonitrile; mobile phase C: methanol) were employed to run a gradient condition from 100 % A to 50% B and 50% C in 10 min., to 100 % B in 1 min, 100% B for 3 min. and reequilibrate with 100 % A for 2.5 min)

\*2: HCl salt

\*3: Crystallized on standing from the compound purified (silica gel column chromatography) and dried.

#### Test Example [CRF receptor binding test]

Monkey amygdala membranes were used as a receptor preparation.

<sup>125</sup>I-CRF was used as <sup>125</sup>I-labeled ligand.

Binding reaction using the <sup>125</sup>I-labeled ligand was carried out by the following method described in The Journal of Neuroscience, 7, 88 (1987).

#### Preparation of receptor membranes:

Monkey amygdala was homogenized in 50 mM Tris-HCl buffer (pH 7.0) containing 10 mM MgCl<sub>2</sub>, 2 mM EDTA and centrifuged at 48,000 x g for 20 min, and the precipitate was washed once with Tris-HCl buffer. The washed precipitate was suspended in 50 mM Tris-HCl buffer (pH 7.0) containing 10 mM MgCl<sub>2</sub>, 2 mM EDTA, 0.1% bovine serum albumin and 100 kallikrein units/ml aprotinin, to obtain a membrane preparation.

CRF receptor binding test:

The membrane preparation (0.3 mg protein/ml),  $^{125}\text{I}$ -CRF (0.2 nM) and a test drug were reacted at 25°C for 2 hours. After completion of the reaction, the reaction mixture was filtered by suction through a glass filter (GF/C) treated with  
5 0.3% polyethylene imine, and the glass filter was washed three times with phosphate-buffered saline containing 0.01% Triton X-100. After the washing, the radioactivity of the filter paper was measured in a gamma counter.

The amount of  $^{125}\text{I}$ -CRF bound when the reaction was carried out in the presence of 1  $\mu\text{M}$  CRF was taken as the degree of nonspecific binding of  $^{125}\text{I}$ -CRF,  
10 and the difference between the total degree of  $^{125}\text{I}$ -CRF binding and the degree of nonspecific  $^{125}\text{I}$ -CRF binding was taken as the degree of specific  $^{125}\text{I}$ -CRF binding. An inhibition curve was obtained by reacting a definite concentration (0.2 nM) of  $^{125}\text{I}$ -CRF with various concentrations of each test drug under the conditions described above. A concentration of the test drug at which binding of  $^{125}\text{I}$ -CRF is  
15 inhibited by 50% ( $\text{IC}_{50}$ ) was determined from the inhibition curve.

As a result, it was found that compounds 1-003, 1-004, 1-006, 1-010, 1-012, 1-013, 1-014, 1-015, 1-016, 1-017, 1-018, 1-019, 1-020, 1-021, 1-024, 1-025, 1-026, 1-027 and 1-028 can be exemplified as typical compounds having an  $\text{IC}_{50}$  value of 100 nM or less.


20

## EFFECT OF THE INVENTION


According to the present invention, compounds having a high affinity for CRF receptors have been provided. These compounds are effective against diseases in which CRF is considered to be involved, such as depression,  
25 anxiety, Alzheimer's disease, Parkinson's disease, Huntington's chorea, eating disorder, hypertension, gastral diseases, drug dependence, epilepsy, cerebral infarction, cerebral ischemia, cerebral edema, cephalic external wound, inflammation, immunity-related diseases, alpecia, irritable bowel syndrome, sleep disorders, dermatitides, schizophrenia, pain, etc.

## CLAIMS

1. A thienopyrimidine or thienopyridine derivative substituted with a cyclic amino group represented by the following formula [I]:



(wherein the cyclic amino group is represented by the following formula [II]:



in which the cyclic amino group is a 3- to 8-membered saturated cyclic amine or a 3- to 8-membered saturated cyclic amine bridged with C<sub>1-5</sub>alkylene or C<sub>1-4</sub>alkylene-O-C<sub>1-4</sub>alkylene between any different two carbon atoms of the cyclic amine, which cyclic amine is substituted with a group represented by -(CR<sup>1</sup>R<sup>2</sup>)<sub>m</sub>-(CHR<sup>3</sup>)<sub>n</sub>-X, R<sup>4</sup> and R<sup>5</sup> independently on the same or different carbon atoms of the cyclic amine;

X is cyano, hydroxy, -CO<sub>2</sub>R<sup>8</sup> or -CONR<sup>9</sup>R<sup>10</sup>;

Y is N or CR<sup>11</sup>;

R<sup>1</sup> is hydrogen, hydroxy, C<sub>1-5</sub>alkyl, C<sub>1-5</sub>alkoxy-C<sub>1-5</sub>alkyl or hydroxy-C<sub>1-5</sub>alkyl;

R<sup>2</sup> is hydrogen or C<sub>1-5</sub>alkyl;

R<sup>3</sup> is hydrogen, cyano, C<sub>1-5</sub>alkyl, C<sub>1-5</sub>alkoxy-C<sub>1-5</sub>alkyl or hydroxy-C<sub>1-5</sub>alkyl;

m is an integer selected from 0, 1, 2, 3, 4 and 5;

n is 0 or 1;

R<sup>4</sup> is hydrogen, hydroxy, hydroxy-C<sub>1-5</sub>alkyl, cyano, cyano-C<sub>1-5</sub>alkyl or C<sub>1-5</sub>alkyl;

R<sup>5</sup> is hydrogen or C<sub>1-5</sub>alkyl;

R<sup>6</sup> is hydrogen, C<sub>1-5</sub>alkyl, C<sub>3-8</sub>cycloalkyl, C<sub>3-8</sub>cycloalkyl-C<sub>1-5</sub>alkyl, hydroxy, C<sub>1-5</sub>alkoxy, C<sub>3-8</sub>cycloalkyloxy, halogen, C<sub>1-5</sub>alkylthio or -N(R<sup>12</sup>)R<sup>13</sup>;

$R^7$  is hydrogen, halogen,  $C_{1-5}$ alkyl,  $C_{3-8}$ cycloalkyl,  $C_{3-8}$ cycloalkyl- $C_{1-5}$ alkyl, hydroxy,  $C_{1-5}$ alkoxy,  $C_{3-8}$ cycloalkyloxy,  $-N(R^{14})R^{15}$ ,  $-CO_2R^{16}$ ,  $-CON(R^{17})R^{18}$ , cyano, nitro,  $C_{1-5}$ alkylthio, trifluoromethyl or trifluoromethoxy;

Ar is aryl or heteroaryl which aryl or heteroaryl is unsubstituted or substituted with 1 or more substituents, which are the same or different, selected from the group consisting of halogen,  $C_{1-5}$ alkyl,  $C_{3-8}$ cycloalkyl,  $C_{2-5}$ alkenyl,  $C_{2-5}$ alkynyl,  $C_{1-5}$ alkoxy,  $C_{1-5}$ alkylthio,  $C_{1-5}$ alkylsulfinyl,  $C_{1-5}$ alkylsulfonyl, cyano, nitro, hydroxy,  $-CO_2R^{19}$ ,  $-C(=O)R^{20}$ ,  $-CONR^{21}R^{22}$ ,  $-OC(=O)R^{23}$ ,  $-NR^{24}CO_2R^{25}$ ,  $-S(=O)_rNR^{26}R^{27}$ , trifluoromethyl, trifluoromethoxy, difluoromethoxy, fluoromethoxy, methylenedioxy, ethylenedioxy and  $-N(R^{28})R^{29}$ ;

$R^8$  is hydrogen,  $C_{1-10}$ alkyl,  $C_{3-8}$ cycloalkyl,  $C_{3-8}$ cycloalkyl- $C_{1-5}$ alkyl, aryl or aryl- $C_{1-5}$ alkyl;

$R^9$  and  $R^{10}$  are the same or different, and independently are hydrogen,  $C_{1-5}$ alkyl,  $C_{3-8}$ cycloalkyl,  $C_{3-8}$ cycloalkyl- $C_{1-5}$ alkyl, aryl or aryl- $C_{1-5}$ alkyl; or  $R^9$  and  $R^{10}$  form a ring selected from saturated 3 to 8 membered ring with the attached nitrogen atom, wherein one of the carbon atoms of such saturated 3 to 8 membered ring is optionally replaced by an oxygen or sulfur atom or by N-Z wherein Z is hydrogen, benzyl or  $C_{1-5}$ alkyl;

$R^{11}$  is hydrogen, halogen or  $C_{1-5}$ alkyl;

$R^{12}$ ,  $R^{13}$ ,  $R^{14}$  and  $R^{15}$  are the same or different, and independently are hydrogen or  $C_{1-5}$ alkyl;

$R^{16}$ ,  $R^{19}$  and  $R^{25}$  are the same or different, and independently are hydrogen or  $C_{1-5}$ alkyl,  $C_{3-8}$ cycloalkyl,  $C_{3-8}$ cycloalkyl- $C_{1-5}$ alkyl, aryl or aryl- $C_{1-5}$ alkyl;


$R^{17}$ ,  $R^{18}$ ,  $R^{20}$ ,  $R^{21}$ ,  $R^{22}$ ,  $R^{23}$ ,  $R^{24}$ ,  $R^{26}$ ,  $R^{27}$ ,  $R^{28}$  and  $R^{29}$  are the same or different, and independently are hydrogen,  $C_{1-5}$ alkyl or  $C_{3-8}$ cycloalkyl;

r is 1 or 2)

, individual isomers thereof or racemic or non-racemic mixtures of isomers thereof, pharmaceutically acceptable prodrugs thereof or pharmaceutically acceptable salts and hydrates thereof.

2. The thienopyrimidine derivative substituted with the cyclic amino group according to claim 1 represented by the following formula [III]:





(wherein X, m, n, the cyclic amino group,  $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^4$ ,  $R^5$ ,  $R^6$ ,  $R^7$  and Ar are as defined in claim 1), individual isomers thereof or racemic or non-racemic mixtures of isomers thereof, or pharmaceutically acceptable salts and hydrates thereof.

3. The thienopyrimidine derivative substituted with the cyclic amino group according to claim 2 represented by formula [III], wherein X is cyano; the cyclic amino group is a 4- to 7-membered saturated cyclic amine; n is 0; m is an integer selected from 0, 1, 2 and 3;  $R^1$ ,  $R^2$ ,  $R^4$  and  $R^5$  are hydrogen;  $R^6$  is  $C_{1-5}$ alkyl;  $R^7$  is hydrogen or  $C_{1-5}$ alkyl; and Ar is phenyl which phenyl is substituted with two or three substituents, which are the same or different, selected from the group consisting of halogen,  $C_{1-3}$ alkyl,  $C_{1-3}$ alkoxy,  $C_{1-3}$ alkylthio, trifluoromethyl, trifluoromethoxy and  $-N(R^{28})R^{29}$  (wherein  $R^{28}$  and  $R^{29}$  are the same or different, and independently are hydrogen or  $C_{1-3}$ alkyl), individual isomers thereof or racemic or non-racemic mixtures of isomers thereof, or pharmaceutically acceptable salts and hydrates thereof.

4. The thienopyrimidine derivative substituted with the cyclic amino group according to claim 2 represented by formula [III], wherein X is cyano; the cyclic amino group is a 6-membered saturated cyclic amine; n is 0; m is 0 or 1;  $R^1$ ,  $R^2$ ,  $R^4$  and  $R^5$  are hydrogen;  $R^6$  is  $C_{1-5}$ alkyl;  $R^7$  is hydrogen or  $C_{1-5}$ alkyl; and Ar is phenyl which phenyl is substituted with two or three substituents, which are the same or different, selected from the group consisting of halogen and  $C_{1-3}$ alkyl, individual isomers thereof or racemic or non-racemic mixtures of isomers thereof, or pharmaceutically acceptable salts and hydrates thereof.

5. The thienopyrimidine derivative substituted with the cyclic amino group


according to claim 2 represented by formula [III], wherein X is hydroxy; the cyclic amino group is a 4- to 7-membered saturated cyclic amine; n is 0; m is an integer selected from 1, 2 and 3;  $R^1$ ,  $R^2$ ,  $R^4$  and  $R^5$  are hydrogen;  $R^6$  is  $C_{1-5}$ alkyl;  $R^7$  is hydrogen or  $C_{1-5}$ alkyl; and Ar is phenyl which phenyl is substituted with two or three substituents, which are the same or different, selected from the group consisting of halogen,  $C_{1-3}$ alkyl,  $C_{1-3}$ alkoxy,  $C_{1-3}$ alkylthio, trifluoromethyl, trifluoromethoxy and  $-N(R^{28})R^{29}$  (wherein  $R^{28}$  and  $R^{29}$  are the same or different, and independently are hydrogen or  $C_{1-3}$ alkyl), individual isomers thereof or racemic or non-racemic mixtures of isomers thereof, or pharmaceutically acceptable salts and hydrates thereof.

6. The thienopyrimidine derivative substituted with the cyclic amino group according to claim 2 represented by formula [III], wherein X is hydroxy; the cyclic amino group is a 6-membered saturated cyclic amine; n is 0; m is an integer selected from 1, 2 and 3;  $R^1$ ,  $R^2$ ,  $R^4$  and  $R^5$  are hydrogen;  $R^6$  is  $C_{1-5}$ alkyl;  $R^7$  is hydrogen or  $C_{1-5}$ alkyl; and Ar is phenyl which phenyl is substituted with two or three substituents, which are the same or different, selected from the group consisting of halogen and  $C_{1-3}$ alkyl, individual isomers thereof or racemic or non-racemic mixtures of isomers thereof, or pharmaceutically acceptable salts and hydrates thereof.

7. The thienopyrimidine derivative substituted with the cyclic amino group according to claim 2 represented by formula [III], wherein X is  $-CO_2R^8$  or  $-CONR^9R^{10}$ ; the cyclic amino group is a 4- to 7-membered saturated cyclic amine; n is 0; m is an integer selected from 0, 1, 2 and 3;  $R^1$ ,  $R^2$ ,  $R^4$  and  $R^5$  are hydrogen;  $R^6$  is  $C_{1-5}$ alkyl;  $R^7$  is hydrogen or  $C_{1-5}$ alkyl;  $R^8$  is hydrogen or  $C_{1-10}$ alkyl;  $R^9$  and  $R^{10}$  are the same or different, and independently are hydrogen or  $C_{1-5}$ alkyl; and Ar is phenyl which phenyl is substituted with two or three substituents, which are the same or different, selected from the group consisting of halogen,  $C_{1-3}$ alkyl,  $C_{1-3}$ alkoxy,  $C_{1-3}$ alkylthio, trifluoromethyl, trifluoromethoxy and  $-N(R^{28})R^{29}$  (wherein  $R^{28}$  and  $R^{29}$  are the same or different, and independently are hydrogen or  $C_{1-3}$ alkyl), individual isomers thereof or racemic or non-racemic mixtures of isomers thereof, or pharmaceutically acceptable salts and hydrates thereof.

8. The thienopyrimidine derivative substituted with the cyclic amino group according to claim 2 represented by formula [III], wherein X is  $-\text{CO}_2\text{R}^8$  or  $-\text{CONR}^9\text{R}^{10}$ ; the cyclic amino group is a 6-membered saturated cyclic amine; n is 0; m is an integer selected from 0, 1, 2 and 3;  $\text{R}^1$ ,  $\text{R}^2$ ,  $\text{R}^4$  and  $\text{R}^5$  are hydrogen;  $\text{R}^6$  is  $\text{C}_{1-5}$ alkyl;  $\text{R}^7$  is hydrogen or  $\text{C}_{1-5}$ alkyl;  $\text{R}^8$  is hydrogen or  $\text{C}_{1-10}$ alkyl;  $\text{R}^9$  and  $\text{R}^{10}$  are the same or different, and independently are hydrogen or  $\text{C}_{1-5}$ alkyl; and Ar is phenyl which phenyl is substituted with two or three substituents, which are the same or different, selected from the group consisting of halogen and  $\text{C}_{1-3}$ alkyl, individual isomers thereof or racemic or non-racemic mixtures of isomers thereof, or pharmaceutically acceptable salts and hydrates thereof.

9. The thienopyridine derivative substituted with the cyclic amino group according to claim 1 represented by the following formula [IV]:



(wherein X, m, n, the cyclic amino group,  $\text{R}^1$ ,  $\text{R}^2$ ,  $\text{R}^3$ ,  $\text{R}^4$ ,  $\text{R}^5$ ,  $\text{R}^6$ ,  $\text{R}^7$ ,  $\text{R}^{11}$  and Ar are as defined in claim 1), individual isomers thereof or racemic or non-racemic mixtures of isomers thereof, or pharmaceutically acceptable salts and hydrates thereof.

10. The thienopyridine derivative substituted with the cyclic amino group according to claim 9 represented by formula [IV], wherein X is cyano; the cyclic amino group is a 4- to 7-membered saturated cyclic amine; n is 0; m is an integer selected from 1, 2 and 3;  $\text{R}^1$ ,  $\text{R}^2$ ,  $\text{R}^4$  and  $\text{R}^5$  are hydrogen;  $\text{R}^6$  is  $\text{C}_{1-5}$ alkyl;  $\text{R}^7$  is hydrogen or  $\text{C}_{1-5}$ alkyl;  $\text{R}^{11}$  is hydrogen or  $\text{C}_{1-5}$ alkyl; and Ar is phenyl which phenyl is substituted with two or three substituents, which are the same or different, selected from the group consisting of halogen,  $\text{C}_{1-3}$ alkyl,  $\text{C}_{1-3}$ alkoxy,  $\text{C}_{1-3}$ alkylthio, trifluoromethyl, trifluoromethoxy and  $-\text{N}(\text{R}^{28})\text{R}^{29}$  (wherein  $\text{R}^{28}$  and  $\text{R}^{29}$  are the same or different, and independently are hydrogen or  $\text{C}_{1-3}$ alkyl), individual isomers

thereof or racemic or non-racemic mixtures of isomers thereof, or pharmaceutically acceptable salts and hydrates thereof.

11. The thienopyridine derivative substituted with the cyclic amino group according to claim 9 represented by formula [IV], wherein X is cyano; the cyclic amino group is a 6-membered saturated cyclic amine; n is 0; m is 0 or 1;  $R^1$ ,  $R^2$ ,  $R^4$  and  $R^5$  are hydrogen;  $R^6$  is  $C_{1-5}$ alkyl;  $R^7$  is hydrogen or  $C_{1-5}$ alkyl;  $R^{11}$  is hydrogen; and Ar is phenyl which phenyl is substituted with two or three substituents, which are the same or different, selected from the group consisting of halogen and  $C_{1-3}$ alkyl, individual isomers thereof or racemic or non-racemic mixtures of isomers thereof, or pharmaceutically acceptable salts and hydrates thereof.

12. The thienopyridine derivative substituted with the cyclic amino group according to claim 9 represented by formula [IV], wherein X is hydroxy; the cyclic amino group is a 4- to 7-membered saturated cyclic amine; n is 0; m is an integer selected from 1, 2 and 3;  $R^1$ ,  $R^2$ ,  $R^4$  and  $R^5$  are hydrogen;  $R^6$  is  $C_{1-5}$ alkyl;  $R^7$  is hydrogen or  $C_{1-5}$ alkyl;  $R^{11}$  is hydrogen or  $C_{1-5}$ alkyl; and Ar is phenyl which phenyl is substituted with two or three substituents, which are the same or different, selected from the group consisting of halogen,  $C_{1-3}$ alkyl,  $C_{1-3}$ alkoxy,  $C_{1-3}$ alkylthio, trifluoromethyl, trifluoromethoxy and  $-N(R^{28})R^{29}$  (wherein  $R^{28}$  and  $R^{29}$  are the same or different, and independently are hydrogen or  $C_{1-3}$ alkyl), individual isomers thereof or racemic or non-racemic mixtures of isomers thereof, or pharmaceutically acceptable salts and hydrates thereof.

13. The thienopyridine derivative substituted with the cyclic amino group according to claim 9 represented by formula [IV], wherein X is hydroxy; the cyclic amino group is a 6-membered saturated cyclic amine; n is 0; m is an integer selected from 1, 2 and 3;  $R^1$ ,  $R^2$ ,  $R^4$  and  $R^5$  are hydrogen;  $R^6$  is  $C_{1-5}$ alkyl;  $R^7$  is hydrogen or  $C_{1-5}$ alkyl;  $R^{11}$  is hydrogen; and Ar is phenyl which phenyl is substituted with two or three substituents, which are the same or different, selected from the group consisting of halogen and  $C_{1-3}$ alkyl, individual isomers thereof or racemic or non-racemic mixtures of isomers thereof, or pharmaceutically acceptable salts and hydrates thereof.

14. The thienopyridine derivative substituted with the cyclic amino group according to claim 9 represented by formula [IV], wherein X is  $-\text{CO}_2\text{R}^8$  or  $-\text{CONR}^9\text{R}^{10}$ ; the cyclic amino group is a 4- to 7-membered saturated cyclic amine; n is 0; m is an integer selected from 0, 1, 2 and 3;  $\text{R}^1$ ,  $\text{R}^2$ ,  $\text{R}^4$  and  $\text{R}^5$  are hydrogen;  $\text{R}^6$  is  $\text{C}_{1-5}$ alkyl;  $\text{R}^7$  is hydrogen or  $\text{C}_{1-5}$ alkyl;  $\text{R}^8$  is hydrogen or  $\text{C}_{1-10}$ alkyl;  $\text{R}^9$  and  $\text{R}^{10}$  are the same or different, and independently are hydrogen or  $\text{C}_{1-5}$ alkyl;  $\text{R}^{11}$  is hydrogen or  $\text{C}_{1-5}$ alkyl; and Ar is phenyl which phenyl is substituted with two or three substituents, which are the same or different, selected from the group consisting of halogen,  $\text{C}_{1-3}$ alkyl,  $\text{C}_{1-3}$ alkoxy,  $\text{C}_{1-3}$ alkylthio, trifluoromethyl, trifluoromethoxy and  $-\text{N}(\text{R}^{28})\text{R}^{29}$  (wherein  $\text{R}^{28}$  and  $\text{R}^{29}$  are the same or different, and independently are hydrogen or  $\text{C}_{1-3}$ alkyl), individual isomers thereof or racemic or non-racemic mixtures of isomers thereof, or pharmaceutically acceptable salts and hydrates thereof.

15. The thienopyridine derivative substituted with the cyclic amino group according to claim 2 represented by formula [IV], wherein X is  $-\text{CO}_2\text{R}^8$  or  $-\text{CONR}^9\text{R}^{10}$ ; the cyclic amino group is a 6-membered saturated cyclic amine; n is 0; m is an integer selected from 0, 1, 2 and 3;  $\text{R}^1$ ,  $\text{R}^2$ ,  $\text{R}^4$  and  $\text{R}^5$  are hydrogen;  $\text{R}^6$  is  $\text{C}_{1-5}$ alkyl;  $\text{R}^7$  is hydrogen or  $\text{C}_{1-5}$ alkyl;  $\text{R}^8$  is hydrogen or  $\text{C}_{1-10}$ alkyl;  $\text{R}^9$  and  $\text{R}^{10}$  are the same or different, and independently are hydrogen or  $\text{C}_{1-5}$ alkyl;  $\text{R}^{11}$  is hydrogen; and Ar is phenyl which phenyl is substituted with two or three substituents, which are the same or different, selected from the group consisting of halogen and  $\text{C}_{1-3}$ alkyl, individual isomers thereof or racemic or non-racemic mixtures of isomers thereof, or pharmaceutically acceptable salts and hydrates thereof.

16. Compounds represented by formula [I] according to claim 1, which compounds are selected from the group consisting of

{1-[7-(4-Bromo-2,6-dimethyl-phenyl)-2-methyl-thieno[3,2-d]pyrimidin-4-yl]-piperidin-4-yl}-methanol,

{1-[7-(4-bromo-2,6-dimethyl-phenyl)-2,6-dimethyl-thieno[3,2-d]pyrimidin-4-yl]-piperidin-4-yl}-methanol,

2-{1-[7-(4-bromo-2,6-dimethyl-phenyl)-2,6-dimethyl-thieno[3,2-d]pyrimidin-4-yl]-piperidin-4-yl}-ethanol,  
    {1-[7-(4-bromo-2,6-dimethyl-phenyl)-2,6-dimethyl-thieno[3,2-d]pyrimidin-4-yl]-piperidin-4-yl}-acetonitrile,  
    {1-[3-(2,4-dichloro-phenyl)-5-methyl-thieno[3,2-b]pyridin-7-yl]-piperidin-4-yl}-methanol,  
    {1-[5-methyl-3-(2,4,6-trimethyl-phenyl)-thieno[3,2-b]pyridin-7-yl]-piperidin-4-yl}-methanol,  
    {1-[3-(4-bromo-2,6-dimethyl-phenyl)-5-methyl-thieno[3,2-b]pyridin-7-yl]-piperidin-4-yl}-methanol,  
    {1-[3-(4-bromo-2,6-dimethyl-phenyl)-2,5-dimethyl-thieno[3,2-b]pyridin-7-yl]-piperidin-4-yl}-methanol,  
    {1-[3-(2,4-dibromo-phenyl)-5-methyl-thieno[3,2-b]pyridin-7-yl]-piperidin-4-yl}-methanol,  
    {1-[5-methyl-3-(2,4,6-trichloro-phenyl)-thieno[3,2-b]pyridin-7-yl]-piperidin-4-yl}-methanol,  
2-{1-[3-(4-bromo-2,6-dimethyl-phenyl)-5-methyl-thieno[3,2-b]pyridin-7-yl]-piperidin-4-yl}-ethanol,  
2-{1-[3-(4-bromo-2,6-dimethyl-phenyl)-2,5-dimethyl-thieno[3,2-b]pyridin-7-yl]-piperidin-4-yl}-ethanol,  
2-{1-[3-(2,4-dibromo-phenyl)-5-methyl-thieno[3,2-b]pyridin-7-yl]-piperidin-4-yl}-ethanol,  
2-{1-[5-methyl-3-(2,4,6-trichloro-phenyl)-thieno[3,2-b]pyridin-7-yl]-piperidin-4-yl}-ethanol,  
1-[5-methyl-3-(2,4,6-trimethyl-phenyl)-thieno[3,2-b]pyridin-7-yl]-piperidine-3-carbonitrile,  
    {1-[3-(4-bromo-2,6-dimethyl-phenyl)-5-methyl-thieno[3,2-b]pyridin-7-yl]-piperidin-4-yl}-acetonitrile,  
    {1-[3-(4-bromo-2,6-dimethyl-phenyl)-2,5-dimethyl-thieno[3,2-b]pyridin-7-yl]-piperidin-4-yl}-acetonitrile,  
    {1-[3-(2,4-dibromo-phenyl)-5-methyl-thieno[3,2-b]pyridin-7-yl]-piperidin-4-yl}-acetonitrile  
and {1-[5-methyl-3-(2,4,6-trichloro-phenyl)-thieno[3,2-b]pyridin-7-yl]-

piperidin-4-yl}-acetonitrile.

17. An antagonist for CRF receptors, comprising a thienopyrimidine or thienopyridine derivative substituted with a cyclic amino group, a pharmaceutically acceptable salt thereof or its hydrate according to any one of claims 1 to 16, as an active ingredient.

18. Use of a thienopyrimidine or thienopyridine derivative substituted with a cyclic amino group, a pharmaceutically acceptable salt thereof or its hydrate according to any one of claim 1 to 16, for the manufacture of a therapeutic agent as an antagonist for CRF receptors.

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/JP2005/000318

A. CLASSIFICATION OF SUBJECT MATTER  
IPC 7 C07D495/04

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data, BEILSTEIN Data, PAJ, WPI Data

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 02/02549 A (TAISHO PHARMACEUTICAL CO., LTD; NAKAZATO, ATSURO; KUMAGAI, TOSHIHITO;) 10 January 2002 (2002-01-10) cited in the application page 1, line 2 - line 10 page 4, line 9 - line 10 page 5, form (11) page 78, line 19 - page 79, line 2 -----	1-18
A	WO 97/29110 A (JANSSEN PHARMACEUTICA N.V; NEUROCRINE BIOSCIENCES INC; CHEN, CHEN; WEB) 14 August 1997 (1997-08-14) cited in the application page 1, line 5 - line 8 page 3, Formula (I) Table 2, Co. No. 36-40 Table 3, Co. No. 59, 60 page 11, line 32 - page 12, line 2 ----- -/--	1-18



Further documents are listed in the continuation of box C.



Patent family members are listed in annex

\* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance, the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

21 June 2005

Date of mailing of the international search report

28/06/2005

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx 31 651 epo nl,  
Fax (+31-70) 340-3016

Authorized officer

Hoepfner, W



## INTERNATIONAL SEARCH REPORT

International Application No

PCT/JP2005/000318

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 98/47903 A (JANSSEN PHARMACEUTICA N.V; NEUROCRINE BIOSCIENCES INC; WEBB, THOMAS, R) 29 October 1998 (1998-10-29) cited in the application page 1, line 5 - line 8 page 3, Formula (I) page 11, line 36 - page 12, line 5 -----	1-18

## INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/JP2005/000318

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0202549	A	10-01-2002	AU 6943701 A	14-01-2002
			BG 107374 A	30-09-2004
			BR 0112166 A	02-09-2003
			CA 2412287 A1	10-01-2002
			CN 1439001 A	27-08-2003
			CN 1535968 A	13-10-2004
			CZ 20024229 A3	18-06-2003
			EA 5289 B1	30-12-2004
			EE 200300007 A	16-08-2004
			EP 1299378 A1	09-04-2003
			HU 0301165 A2	28-08-2003
			WO 0202549 A1	10-01-2002
			JP 2004502685 T	29-01-2004
			MX PA02012820 A	14-05-2003
			NO 20026125 A	04-02-2003
			PL 358411 A1	09-08-2004
			SK 132003 A3	05-08-2003
			TW 591022 B	11-06-2004
			US 2004034061 A1	19-02-2004
			US 2005009874 A1	13-01-2005
			ZA 200210041 A	11-12-2003
WO 9729110	A	14-08-1997	AT 208395 T	15-11-2001
			AU 725674 B2	19-10-2000
			AU 1720997 A	28-08-1997
			CA 2233307 A1	14-08-1997
			DE 69708059 D1	13-12-2001
			DE 69708059 T2	18-07-2002
			DK 882051 T3	25-02-2002
			WO 9729110 A1	14-08-1997
			EP 0882051 A1	09-12-1998
			ES 2167710 T3	16-05-2002
			ID 15904 A	14-08-1997
			JP 2000504678 T	18-04-2000
			NO 981356 A	09-07-1998
			NZ 330118 A	27-03-2000
			PT 882051 T	29-04-2002
			TW 448178 B	01-08-2001
			US 6255310 B1	03-07-2001
			US 2002052362 A1	02-05-2002
			ZA 9700988 A	07-09-1998
WO 9847903	A	29-10-1998	AU 751710 B2	22-08-2002
			AU 7526898 A	13-11-1998
			CA 2272292 A1	29-10-1998
			WO 9847903 A1	29-10-1998
			EP 0977759 A1	09-02-2000
			JP 2002501493 T	15-01-2002
			NZ 335823 A	29-06-2001
			TW 495511 B	21-07-2002
			US 6211195 B1	03-04-2001
			ZA 9803351 A	21-10-1999